


Maricopa County Department of Public Health INFLUENZA UPDATE

October 2, 2011 through March 10, 2012 Posted March 15, 2012

Individuals with Laboratory Confirmed Influenza

This influenza season (2011-12), there have been 294 individuals confirmed with an influenza test. The chart below shows tests each week for the current season (red) and previous seasons. The broken line represents weeks for which all test results have not yet been collected.



Types of Influenza

This influenza season, three types of influenza have been reported. Table 1 shows the number and percentage of tests reported by influenza type.

Influenza Type	Number of Tests	Percentage
Swine flu (Influenza A, type H1N1)	126	63.0%
Seasonal influenza (Influenza A, type H3)	61	30.5%
Influenza B, all types	13	6.5%
		100%
No type determined	94	--
Total	294	--


Pneumonia and Influenza (P&I) Deaths

This influenza season, there have been 599 pneumonia and influenza (P&I) deaths (including 4 pediatric deaths). This is down from last season's total of 699 P&I deaths (including 11 pediatric deaths). Table 2 shows the number of P&I deaths and the number of influenza deaths that occurred this season and last season.


	Current Season (2011-2012) As of [3/10/2012]		Last Season (2010-2011) As of [3/12/2011]	
	Pneumonia and Influenza	Influenza	Pneumonia and Influenza	Influenza
Pediatric (Under age 18)	4	0	11	3
Adult (18 and Over)	595	1	688	9

Respiratory Illness Activity at Hospital Emergency Departments

This influenza season, there have been 7,812 visits to emergency departments by individuals with flu-like symptoms. To be counted as an individual with flu-like symptoms, the patient must have both fever and cough or sore throat. Graph 2 shows the number of patients with flu-like symptoms seen at 18 emergency departments. The broken bars represent weeks for which all visits have not yet been collected.




Graph 3 shows what percent of all emergency department visits were individuals with flu-like symptoms (fever and cough/sore throat). The graph shows the trend week by week, with the current season in red. The broken line represents weeks for which all visits have not been collected.



Respiratory Illness Activity at Schools

Graph 4 shows the percent of enrolled students in selected public schools who were absent due to flu-like symptoms. To be counted as a student absent with flu-like symptoms, the student must be kept home from school and reported (usually by a parent) to have both a fever and cough or sore throat. The chart shows the trend week by week, for the current school year (red) and previous school years.



*The Maricopa County Department of Public Health wishes to thank all of the
influenza surveillance participants:*

*Arizona Department of Health Services
Local Hospital Emergency Departments & Infection Control Practitioners
(CDC's Biosense Influenza Module, 12 additional hospitals not a part of the Influenza Module)
Maricopa County Public Schools
Maricopa County Office of Vital Statistics*

Questions about this report may be directed to:

epidemiology@mail.maricopa.gov

**Information on influenza activity for the state, the U.S. and the world may be found
at the following websites:**

<http://www.azdhs.gov/phs/oids/epi/flu/index.htm> (ADHS)

<http://www.cdc.gov/flu/weekly/fluactivitysurv.htm> (CDC)

http://www.who.int/influenza/gisrs_laboratory/updates/en/index.html (WHO)