


Maricopa County Department of Public Health INFLUENZA UPDATE

September 30, 2012 through March 30, 2013

Posted April 4, 2013

Individuals with Laboratory Confirmed Influenza

This influenza season (2012-13), there have been 6,887 individuals confirmed with an influenza test, 84 cases reported within the past week. On October 30th, 2012, the first locally-acquired influenza case was confirmed at the Arizona State Public Health Laboratory. The state of Arizona is reporting **local** activity. The numbers in this report are based on processed cases. There are a large number of reports that are still being processed. This means the numbers in this report are **provisional**. The chart below shows tests each week for the current season (red) and previous seasons.



Types of Influenza

This influenza season, three types of influenza have been reported. Table 1 shows the number and percentage of tests reported by influenza type and subtype.

Total	6887	100.0%
Type A	4958	72.0%
<i>Subtype 2009 H1N1</i>	7	0.1%
<i>Subtype H3</i>	285	5.7%
<i>Subtyping not performed or unknown</i>	4666	94.1%
Type B	1589	23.1%
Type Unknown	340	4.9%


Pneumonia and Influenza (P&I) Deaths

This influenza season, there have been 817 pneumonia and influenza (P&I) deaths (including 7 pediatric deaths (none attributed to influenza)). There have been 25 adult influenza deaths reported this season. This has increased from last season's total of 704 P&I deaths (including 6 pediatric deaths (none attributed to influenza)). Table 2 shows the number of P&I deaths and the number of influenza deaths that occurred this season and last season.


	Current Season (2012-2013) As of [03/30/2013]		Last Season (2011-2012) As of [03/31/2012]	
	Pneumonia and Influenza	Influenza	Pneumonia and Influenza	Influenza
Pediatric (Under Age 18)	7	0	6	0
Adult (18 and Over)	810	25	698	4

Respiratory Illness Activity at Hospital Emergency Departments

This influenza season, there have been 9,831 visits to emergency departments by individuals with flu-like symptoms. To be counted as an individual with flu-like symptoms, the patient must have both fever and cough or sore throat. Graph 2 shows the number of patients with flu-like symptoms seen at 12 emergency departments.




Graph 3 shows what percent of all emergency department visits were individuals with flu-like symptoms (fever and cough/sore throat). The graph shows the trend week by week, with the current season in red.



Respiratory Illness Activity at Schools

Graph 4 shows the percent of enrolled students in selected public schools who were absent due to flu-like symptoms. To be counted as a student absent with flu-like symptoms, the student must be kept home from school and reported (usually by a parent) to have both a fever and cough or sore throat. The chart shows the trend week by week, for the current school year (red) and previous school years.



***The Maricopa County Department of Public Health wishes to thank all of the
influenza surveillance participants:***

*Arizona Department of Health Services
Local Hospital Emergency Departments & Infection Control Practitioners
Maricopa County Public Schools
Maricopa County Office of Vital Statistics*

Questions about this report may be directed to:

epidemiology@mail.maricopa.gov

***Information on influenza activity for the state, the U.S. and the world may be found
at the following websites:***

<http://www.azdhs.gov/phs/oids/epi/flu/index.htm> (ADHS)

<http://www.cdc.gov/flu/weekly/fluactivitysurv.htm> (CDC)

http://www.who.int/influenza/gisrs_laboratory/updates/en/index.html (WHO)