

Maricopa County Department of Public Health


INFLUENZA UPDATE

September 30, 2012 through July 13, 2013

Posted July 18, 2013

Individuals with Laboratory Confirmed Influenza

This influenza season (2012-13), there have been 6,582 individuals confirmed with an influenza test; 1 new case has been reported in the past two weeks. On October 30th, 2012, the first locally-acquired influenza case was confirmed at the Arizona State Public Health Laboratory. The state of Arizona reported **sporadic** activity for week 27 and **no** activity for week 28. The numbers in this report are based on processed cases. There are a large number of reports that are still being processed. This means the numbers in this report are **provisional**. The chart below shows tests each week for the current season (red) and previous seasons.



Types of Influenza

This influenza season, three types of influenza have been reported. Table 1 shows the number and percentage of tests reported by influenza type and subtype.

Total	6582	100.0%
Type A	4756	72.3%
<i>Subtype 2009 H1N1</i>	30	0.6%
<i>Subtype H3</i>	892	18.8%
<i>Subtyping not performed or unknown</i>	3834	80.6%
Type B	1676	25.5%
Type Unknown	150	2.3%


Pneumonia and Influenza (P&I) Deaths

This influenza season, there have been 1,129 pneumonia and influenza (P&I) deaths (including 11 pediatric deaths (none attributed to influenza)). There have been 26 adult influenza deaths reported this season. This has increased from last season's total of 1,063 P&I deaths (including 9 pediatric deaths (none attributed to influenza)). Table 2 shows the number of P&I deaths and the number of influenza deaths that occurred this season and last season.


	Current Season (2012-2013) As of [07/13/2013]		Last Season (2011-2012) As of [07/14/2012]	
	Pneumonia and Influenza	Influenza	Pneumonia and Influenza	Influenza
Pediatric (Under Age 18)	11	0	9	0
Adult (18 and Over)	1118	26	1054	5

Respiratory Illness Activity at Hospital Emergency Departments

This influenza season, there have been 11,203 visits to emergency departments by individuals with flu-like symptoms. To be counted as an individual with flu-like symptoms, the patient must have both fever and cough or sore throat. Graph 2 shows the number of patients with flu-like symptoms seen at 12 emergency departments.




Graph 3 shows what percent of all emergency department visits were individuals with flu-like symptoms (fever and cough/sore throat). The graph shows the trend week by week, with the current season in red.



Respiratory Illness Activity at Schools

Graph 4 shows the percent of enrolled students in selected public schools who were absent due to flu-like symptoms. To be counted as a student absent with flu-like symptoms, the student must be kept home from school and reported (usually by a parent) to have both a fever and cough or sore throat. The chart shows the trend week by week, for the current school year (red) and previous school years.



***The Maricopa County Department of Public Health wishes to thank all of the
influenza surveillance participants:***

*Arizona Department of Health Services
Local Hospital Emergency Departments & Infection Control Practitioners
Maricopa County Public Schools
Maricopa County Office of Vital Statistics*

***Questions about this report may be directed to:
epidemiology@mail.maricopa.gov***

***Information on influenza activity for the state, the U.S. and the world may be found
at the following websites:***

<http://www.azdhs.gov/phs/oids/epi/flu/index.htm> (ADHS)
<http://www.cdc.gov/flu/weekly/fluactivitysurv.htm> (CDC)
http://www.who.int/influenza/gisrs_laboratory/updates/en/index.html (WHO)